

Turbulence & Phonology

John J. Ohala* & Maria-Josep Solé #

**Department of Linguistics, University of California, Berkeley
ohala@berkeley.edu*

*#Department of English, Universitat Autònoma de Barcelona, Spain
mariajosep.sole@uab.es*

In this paper we aim to provide an account of some of the phonological patterns involving turbulent sounds, summarizing material we have published previously and results from other investigators. In addition, we explore the ways in which sounds pattern, combine, and evolve in language and how these patterns can be derived from a few physical and perceptual principles which are independent from language itself (Lindblom 1984, 1990a) and which can be empirically verified (Ohala and Jaeger 1986). This approach should be contrasted with that of mainstream phonological theory (i.e., phonological theory within generative linguistics) which primarily considers sound structure as motivated by ‘formal’ principles or constraints that are specific to language, rather than relevant to other physical or cognitive domains.

For this reason, the title of this paper is meant to be ambiguous. The primary sense of it refers to sound patterns in languages involving sounds with turbulence, e.g., fricatives and stops bursts, but a secondary meaning is the metaphorical turbulence in the practice of phonology over the past several decades. We’ll treat the latter topic first.

1. Turbulence in Phonology

Anyone familiar with the history of phonological science in the 20th century will have to concede that there has been considerable turbulence in the theoretical domain. To be sure, there were controversies in phonology in the 19th century, too, for example, the dispute as to whether Sanskrit should be taken as the oldest ancestor of what became known as the Indo-European language family or whether an attempt should be made to reconstruct a parent language of which even Sanskrit was an off-shoot. Schleicher, the advocate of the latter view, eventually won that dispute. There were also disputes as to the causes and mechanisms of sound change; these disputes have not been satisfactorily resolved even to this day. Nevertheless, this domain of phonological science -- historical linguistics -- made steady and remarkable progress from its beginning in the 18th century (e.g., ten Kate 1723, des Brosses

1765)¹. The methodology, the “comparative method”, has been refined and proven itself. Beginning around the turn of the 19th to 20th centuries, a new program developed, spurred by the writings of Kruzewski, Baudouin de Courtenay, Saussure, Sapir and others. This was to be an account of the psychological aspect of language, i.e., how individual distinctive speech sounds and their contextual variants arise and are maintained and managed in the mind of the speaker-hearer. Turbulence in linguistic theory arose when patterns arising from physical and physiological factors, that had previously been judged to be extra-linguistic, were claimed by generative grammarians in the mid-20th century to be incorporated into the posited psychological lexicon and grammar of the speaker. The dust still hasn’t settled on this controversy. In form these supposed psychological representations of language and the methods used to discover them were largely identical to the descriptive entities and methods of the historical phonologist.

Against this historical background, we declare that the explanations we give below for sound patterns involving noise (turbulence) are strictly physical phonetic. In our view there is no mystery about how physical phonetic factors can become phonologized and manifested in languages’ sound patterns: the variation in the speech due to these physical constraints can lead to the listener’s misperception, misparsing, and misconstruing of the speaker’s target pronunciation. Listener error, then, can lead to a change in the pronunciation norms just as manuscript copyists’ error led to different variants of ancient texts in the time before printing (Ohala 1981b, 1989).

The ultimate purpose of the experimental approach to phonology illustrated in this paper is to demonstrate that phonological theory must be based on mechanisms and principles coming from the subsystems involved in speech production and speech perception (Ohala 1981a, 1981b, 1992).

For ease of presentation, we group the sound patterns according to aerodynamic factors (section 2), acoustic-auditory factors (section 3), and timing factors (section 4), but most of the patterns involve the interaction between a number of these factors.

2. Aerodynamic factors relevant to turbulence

2.1. Basic principles

From an aerodynamic point of view, we can think of the vocal tract as two air cavities, the lung cavity and the supraglottal cavity, ultimately connected to the atmosphere. The two cavities are connected by the glottis which allows pulmonic air to flow into the oral cavity as

¹ Not to neglect worthy precursors even before that, e.g., van Boixhorn (1647)

pulmonic forces (from muscular activity or passive recoil) compress the lungs. The supraglottal cavity is connected to the atmosphere by the mouth (and the nose) which can impede the air flowing out with changes in articulatory constriction of the lips and tongue. Thus the two main valves (along with the nasal valve) that regulate the airflow used in speech are the glottis and the oral constriction, and turbulent noise can be generated at both of these constrictions.

The generation of audible turbulence, i.e., noise, in the vocal tract is necessary for the production of fricatives, the fricative release of affricates and the burst of stops. However, audible turbulence may also be associated with the production of vowels and sonorants in certain conditions. Although there is some degree of low-level air flow turbulence even for the most open of speech sounds, i.e., something like [ε] (because the air flowing into the vocal tract acquires some turbulence upon passing through the vibrating vocal cords) it is only when the turbulence reaches a level to become audible that it can play some role in speech.

Turbulent airflow is determined by a multiplicity of factors, including roughness and length of the channel, shape of the orifice, and whether the air downstream of the constriction is already turbulent, but it is the *speed* of air flowing through the constriction which is the main factor. The speed of air (or ‘particle velocity’, v), in turn, depends on the volume of air flowing through the constriction (called the ‘volume velocity’, U) and on the cross-sectional area of the constriction (A), as indicated in (1). Thus the larger the volume of air per unit time and the smaller the constriction, the higher the velocity, and the more intense the friction noise.

$$v = U/A \tag{1}$$

where v is the particle velocity in cm/sec; U is the volume velocity in cm³/sec, A is the cross-dimensional area of the constriction in cm².

The volume of air, U , that will flow through a constriction depends on the size of the aperture, A , and the pressure difference across the aperture, that is, the difference between the upstream pressure and downstream pressure, $P_{\text{upstream}} - P_{\text{downstream}}$, as shown in (2) (Warren and DuBois 1964). In the case of an oral constriction, this will be the difference in pressure between the oral cavity and the atmosphere ($P_{\text{oral}} - P_{\text{atmospheric}}$)², and in the case of a glottal

² Although atmospheric pressure (absolute) is roughly 1033 cm H₂O at sea level, it is common to take atmospheric pressure as zero (0) and express pressure in the vocal tract as x cm H₂O with respect to that. Nothing is lost with such a mathematical convention.

constriction, the difference in pressure between the subglottal (or pulmonic) cavity and the oral cavity ($P_{\text{subglottal}} - P_{\text{oral}}$). The greater the difference in pressure and the larger the area of the constriction, the larger the rate of flow. The exponent, a , varies between .5 and 1, depending on the nature of the flow; it is 1 when the flow is smooth or laminar, and .5 when the flow is turbulent. In the conditions found in speech production, i.e., what is called 'nozzle flow', this number may vary continuously between these two extremes (Jaeger and Matthys 1970). Naturally, the direction of air flow will always be from the cavity with greater pressure to that with lesser pressure.

$$U = A (P_{\text{upstream}} - P_{\text{downstream}})^a c \quad \text{or} \quad U = A (\Delta P)^a c \quad (2)$$

where P is the pressure in cm H₂O; c is a constant.

As mentioned, traditionally, the critical velocity at which the change from laminar to turbulent flow occurs is determined by a number of factors, including particle velocity, the diameter and roughness of the channel the air passes through, etc. The relative contribution of these factors for certain flow conditions is quantified in the Reynolds number. When the Reynolds number exceeds a certain threshold the airflow is supposed to change from smooth or 'laminar' to turbulent. However, it is the case that in irregularly-shaped channels like the vocal tract and with airflow that usually has a turbulent entry into the vocal tract (certainly the case as the air passes several "rough" surfaces –narrow alveoli in the lungs, tracheal rings, vocal cords, ventricular folds, epiglottis, etc.), some turbulence, even audible turbulence can occur in conditions where the Reynolds number is far below the ideal threshold between laminar and turbulent flow. So it is simplest just to state the relation between air velocity and noise in a purely qualitative way: the intensity (i.e., loudness) and center frequency (i.e., pitch) of friction noise varies monotonically with the particle velocity of the air flow, as given in (3a), below (Catford 1977: ch. 3; Stevens 1971; Flanagan and Ishizaka 1976; Flanagan, Ishizaka, and Shipley 1975, 1980; Shadle 1990). A variant of this relation, stating that intensity of friction at a supraglottal constriction increases with increasing oral pressure and decreasing aperture of constriction (therefore, conflating principles 1 and 2), is given in (3b) (Stevens 1971).

$$I_{\text{friction}} \sim v \quad (3a)$$

$$I_{\text{friction}} \sim P_o^{3/2} A^{1/2} \quad (3b)$$

The articulatory constriction for vowels, glides and sonorants is not typically narrow enough to cause a pressure difference across the constriction (in other words, P_{oral} is not much higher than $P_{\text{atmospheric}}$), so that particle velocity, v , through the constriction is kept low and does not reach a level sufficient to generate audible friction (but see below for exceptions when these are voiceless). Obstruents such as fricatives, stops and affricates, on the other hand, are produced with a narrow or complete constriction which causes the P_{oral} to rise substantially over $P_{\text{atmospheric}}$; upon release the particle velocity is high and the airflow becomes more turbulent.

Additional turbulence can be also generated when an air jet that has passed through the major oral cavity constriction encounters any sharp discontinuity: either an abrupt enlargement of the channel or the opposite, i.e., an additional barrier or “baffle”. The former occurs (a) when air passes through the vocal cords during voicing (“voicing” consists of periodic short-term noise bursts occurring at a rate equal to the fundamental frequency) and (b) when the air flows past the sharp-edged constriction at the teeth in a labio-dental fricative such as [f]. The latter occurs when an air jet emerging from an apical-alveolar constriction is directed at the upper and lower incisors. (It is this factor which accounts for the somewhat impoverished apical fricatives made by children who have lost their incisors as part of the change from baby teeth to permanent dentition at approximately age 6 and on.)

A final factor needs to be mentioned regarding the acoustic amplitude of the noise which the turbulence generates: other things being equal, the intensity of the noise is greater, the larger is the resonating cavity downstream of the point where the turbulence occurs. For this reason palatal and velar fricatives have more intense noise than labial and labio-dental fricatives.

2.2. Generalizations on phonetic and phonological universals deduced from aerodynamic principles

In the following sections we review a number of sound patterns, involving the emergence or extinction of turbulence, which can be deduced in part from variations in glottal flow, U_g (section 2.2.1), changes in area of oral constriction, A_o (2.2.2), and changes in oral pressure, P_o (2.2.3).

2.2.1. Variations in glottal flow

It is known that the glottis regulates the flow of air from the lungs into the oral cavity. Vibration at the vocal folds for voiced obstruents causes a diminished rate of flow through the glottis and a significantly lower oral pressure vis-à-vis voiceless obstruents which, in contrast, have a large glottal opening and continuous flow. A relatively low oral pressure is necessary to maintain a sufficient pressure differential across the glottis so that there will be continuous transglottal flow and thus voicing during the obstruent³.

Since by principles (2) and (3), intensity of turbulence is dependent on the pressure difference across the oral constriction, a lower oral pressure for voiced obstruents will result in a lower intensity of high frequency noise during the fricative constriction or at stop release vis-à-vis voiceless obstruents. In addition, due to the reduced transglottal flow, voiced obstruents take longer to build up oral pressure behind the oral constriction, which results in a delayed onset of audible friction for fricatives (Solé 2002b) and a weaker burst for stops compared to their voiceless counterparts. Thus, the characteristic cues for obstruency – abrupt amplitude discontinuities and high intensity noise cues – are enhanced in voiceless obstruents due to the larger rate of flow through the glottis. In sum, for aerodynamic and auditory-acoustic reasons *voicelessness favors or enhances obstruency* (i.e., high intensity friction and release burst). The following phonological generalizations can be derived from this principle.

I. Voicelessness favors obstruency

A. Sonorants (glides, laterals and nasals) become fricatives when devoiced.

As stated in 2.1, the common description of the articulatory difference between an approximant and a fricative is that they have different degrees of constriction (e.g., Clark and Yallop 1990:81; Laver 1994: 134-135). A difference based on constriction degree is endorsed by the present structure of the IPA phonetic alphabet. While this is generally an adequate description, there are cases with considerable phonological interest where this is not completely true.

In general, approximants have a constriction that is large enough to allow the airstream to flow through it without causing turbulence. Nevertheless, by equation (1), $v = U/A$, they may cross the threshold into obstruents if the constriction (A) narrows further or if a higher rate of

³ Other factors have been reported to contribute to keeping a low oral pressure for voiced obstruents, for example, an increased volume of the oral cavity (Ewan and Kronen 1972; Kent and Moll, 1969; Bell-Berti 1975) and/or intentional relaxation of vocal tract muscles resulting in more passive expansion of the walls (Svirsky, Stevens, Matthies, Manzella, Perkell, and Wilhelms-Tricarico 1997).

flow (U) passes through the same constriction. Approximants, e.g., [v j w l ɹ ɻ], and nasals, which by definition are non-obstruents, are usually voiced. When voiceless, however, without any variation in the configuration of the oral articulators, they can become fricative (and thus obstruents), e.g., [f ç ɬ φ ʧ ʂ ʃ] and [m̥ n̥ ŋ̥ ɲ̥]. This happens simply due to the increased airflow passing through the constriction created by these consonants -- the increased airflow being caused by the greater opening (and thus lesser resistance to airflow) at the glottis (see Catford 1977:120ff.). Thus, it is the higher rate of flow through the glottis which creates the higher particle velocity of the airflow through the supraglottal constriction (velocity being directly proportional to flowrate, for a given aperture, $v = U/A$) and leads to turbulence. Hence the obstruent character of the voiceless (former) sonorants. Some phonological consequences of the frication of devoiced sonorants are the following.

A.1. Laterals and /r/s

First, there are cases where a fricative and an approximant alternate and there is co-variation between frication and voicelessness, such that the fricative is voiceless and the approximant is voiced. For example, in Kwakiutl (Boas 1947) there are morphophonemic alternations between ‘plain’ (voiceless), ‘hardened’ (ejectives), and ‘weakened’ (voiced) obstruents and laterals. Of special interest is the fact that the lateral alternates also in manner: the plain voiceless is a fricative /ʃ/ and the weakened form is the voiced approximant /l/, as illustrated in (1). In Welsh (Ball and Williams 2000) there are morphophonemic alternants known as ‘soft mutation’, involving the alternation of voiced-voiceless pairs. Part of these alternations involves the alternation of voiceless and voiced laterals and trills, with the voiceless counterpart being fricative [ʃ] and [r̥] (spelled ‘ll’ and ‘rh’, respectively) and the voiced, simple sonorants [l], [r], see example (2).

(1) Kwakiutl (transcription simplified and converted to IPA) (Boas 1947)

/tsʰōʃ/	‘to be black’
/tsʰōlato/	‘black-eared’

(2) Welsh (Ball and Williams 2000)

<i>llyfr</i>	[ʃ]	‘book’
<i>ei lyfr</i>	[l]	‘his book’

A.2 Glides

In Northern and Central Standard Swedish preaspiration of voiceless stops following stressed vowels has commonly been observed. However, fricativization rather than aspiration is produced between long high vowels, which are diphthongal, and the voiceless stop (Millardet 1911; Rositzke 1940; Helgason 2002: 88). No frication, however, is found for the non-high vowels, see (3). Helgason (2002) notes another factor which may contribute to the observed friction: the tendency to produce friction noise at the end of long, close vowels, regardless of whether or not a consonant follows (e.g. *bi* [bi̯] ‘honey bee’; *gud* [gʊ̯βd] ‘God’). Particularly, in a sequence of a long, close vowel and a voiceless stop, the early glottal abduction for the stop (i.e., preaspiration) during the preceding high vowel will enhance the tendency for friction by increasing the velocity of air across the oral constriction. Similar patterns have been reported for the Jutland dialect of Danish by Andersen (1972).

(3) Fricated glides in Swedish (Millardet 1911; Rositzke 1940)

bit [bi̯çt] ‘a bit, a bite’

kut [kʊ̯çt] ‘seal puppy’

BUT:

tack [takh:] ‘thanks’

peka [pehkə] ‘point’

Bauer (1982) reports fricativization of syllable-initial /w/ into either a voiceless bilabial fricative [ɸ] or a voiced labio-dental fricative /v/ in Hong-Kong Cantonese. The voiceless realization—with increased flow of air through the open glottis—may well give rise to turbulence at the oral constriction, without changing the articulatory configuration. However, considering that a voiced realization may also give way to frication, in the pronunciation /v/, we cannot dismiss the possibility that a closer articulatory constriction syllable initially is responsible for the turbulence and spirantization.

Some varieties of American English retain the older pronunciation [ɰ], a labial velar fricative, in words such as *which*, *whether*, and *white*, rather than the voiced approximant [w]. The correspondence of the voiceless fricative and the voiced glide illustrates the covariation between voicing and frication.

A.3 Nasals

In voiced nasals, vocal fold vibration resonates in the oral and nasal cavities resulting in the low frequency resonances and zeroes characteristic of nasals. Voiceless nasals, on the other hand, have an open glottis for most of the oral closure giving rise to turbulence generated primarily at the nostrils (the point of maximum constriction) no matter what the place of articulation of the nasal. Since the turbulence generated at the nostrils is not very intense and is not amplified and shaped by a downstream cavity⁴, the frication has a weak intensity and there will not be much spectral difference between [m̥ n̥ ŋ̥] during the consonant constriction, though of course they can be still be differentiated by their transitions in adjacent vowels. Although Maddieson (1983), reports that spectral differences may be found during the voiceless portion, nevertheless, as noted, place distinctions in voiceless nasals are obscure. Even so, many languages with distinctive voiceless nasals usually have them at more than one place of articulation. These voiceless nasals usually have a brief voiced period in the last portion of the oral closure, thus they are phonetic sequences [ṁn], etc. In this way different places of articulation can be differentiated —both by the distinctive resonances of the voiced nasal and the transitions in adjacent vowels.

Evidence that voiceless nasals are obstruent-like, specifically, fricatives is provided by Ohala & Ohala (1993). First, distinctive voiceless nasals frequently derive from original /s/+nasal clusters. This is the case for Burmese, where present-day /ṅa/ ‘nose’ stems from Proto-Burmese-Loloish *sna (Bradley 1979), and corresponds to orthographic *sna* in Tibetan. Parallel cases are found in Primitive Greek and Old Irish, where /m̥/ and /n̥/ seem to derive from Indo-European *sm, *sn. Second, children learning English sometimes produce target #sm- and #sn- as [m̥] and [n̥], e.g. [m̥æk] *smack* and [n̥id] *sneeze* (references in Ohala & Ohala 1993: 233). These cases suggest that the voiceless nasal is an adequate auditory substitute for a fricative.

B. Emergence of ‘buccal’ fricatives due to /h/ coarticulating with high vowels

A related phonological pattern, due to the interaction of aerodynamic and acoustic factors, is the emergence of supraglottal fricatives when /h/ is coarticulated with high vowels. For example, the glottal fricative /h/ in Japanese has distinct palatal, [ç], and labial, [ɸ], variants before high front and high back vowels, respectively, and [h] before more open vowels. This

⁴ The sound source excites mostly the cavities anterior to the constriction where the sound is generated, whereas the back cavities do not contribute much acoustically (Fant 1960, Stevens 1998). Since the oral and nasal cavities posterior to the nostrils do not contribute resonances, the different nasals do not differ much acoustically.

is illustrated in (4) below. (Note that high vowels are allophonically devoiced between voiceless sounds and optionally devoiced when following a voiceless consonant or utterance-finally):

(4) <i>hikaku</i>	/hikaku/	[ç̥ikaku]	'comparison'
<i>futa</i>	/hʉʉta/	[ç̥ʉʉta]	'lid'
BUT:			
<i>happyaku</i>	/happjaku/	[hap̚ːaku]	'eight hundred'

In present-day English, /h/, which is the voiceless version of the following vowel or glide (Lehiste 1964, ch. 5), has the fricative allophone [ç̥] before /j/ and /i:/, Hugh (a name) [ç̥ju:], heal [ç̥i:t̚]. Similarly, in Fante /h/ is phonetically [ç̥] before /ɪ/ (Schahter and Fromkin 1968). The emergent fricatives in Swedish, illustrated in (3) above, are a further example⁵.

In segments involving a glottal and a supraglottal constriction, and turbulent noise generated at both of these constrictions as is the case for voiceless high vowels and sonorants— acoustic factors are also relevant. In general terms, in sounds with more than one constriction, the major noise source is contributed by the constriction with a smaller cross-sectional area, other things being equal (e.g., smoothness of the channel surface, presence of an obstacle). The area of supraglottal constriction for high vowels like [i] and [u] is about 0.2 to 0.3 cm² (Chiba and Kajiyama 1941; Fant 1960; Baer, Gore, Gracco and Nye 1991), and hence comparable to the area of the open glottis, approximately 0.3cm² (Stevens 1998: 37). Thus, if the two constrictions are of about the same size ($A_o = A_g = 0.3 \text{ cm}^2$) the relative noise level at each constriction is about the same. However, the transfer function of the two sources is not the same, the major noise source is contributed by the outer constriction (Stevens 1998: 442-3). This is because the cavity anterior to the outer constriction enhances the amplitude of the frequencies of the front-cavity resonance for [i] (F3 peak), and thus palatal frication dominates over glottal frication. Similarly, for [ʉ] [u] the amplitude of the labial or velar frication (F2 peak) is greater. Hence the distinct ‘buccal’ fricatives resulting from voiceless high vowels.

C. Stop releases engender frication on adjacent glides, high vowels and sonorants

⁵ A related phenomenon is the preservation of Latin word-initial /f/ only before /w/ in Spanish (e.g., *fuelle* [ˈfwente] < *fonte* ‘fountain’). /f/ became /h/ and was later lost preceding all other vowels (e.g., *hablar* [aˈβlar] < *fabulare* ‘to talk’, *hierro* [ˈjero] < *ferru* ‘iron’, *hondo* [ˈondo] < *fundu* ‘deep’) (Menéndez Pidal 1968: 122).

A stop release is in itself a brief period of turbulence noise – due to the high rate of airflow which arises from the high back pressure developed during the consonantal closure. However in some cases the turbulence noise is prolonged at the release of a stop followed by a high vowel or a glide and this leads to the emergence of a fricative (Stevens 1971, Ohala 1983b) . When a stop is followed by segments involving a high tongue position, such as high vowels and glides (and to a certain extent liquids /r, l/), the air pressure build-up behind the stop constriction is released through a narrow channel (A) which offers a high resistance to exiting air and thus increases the particle velocity and turbulence (by equation (1)). It can take a few tens of milliseconds for the P_{oral} to approach $P_{atmospheric}$ and during this time the air will be forced through the constriction at a higher rate. Hence the initial portions of the vowel or glide can be fricated. The phonologization of the stop as an affricate is due to the listener parsing the prolonged frication with the stop, not the vowel. Such emergent affricates do not develop before more open vowels with a wider constriction.

In addition, onset of vocal fold vibration after a stop is delayed in high vowels and glides vis-à-vis open vowels (Ohala 1976, 1981a, 1983b, Chang 1999), due to the slower release of the oral pressure through the narrow constriction and the longer time needed to achieve the pressure differential for voicing. This results in a longer period of turbulent flow, which contributes to the percept of frication. This is illustrated in Fig. 1 which shows that that oral pressure impulse from the stop decays more slowly for /tja:/, /twa:/, /tra:/ vis-à-vis /ta:/ due to the exiting air encountering greater resistance. For these sequences the high velocity airflow passes through a narrow constriction, enhancing turbulence, for a relatively long time creating the percept of an affricated stop release.

INSERT FIG. 1 HERE

Although most phonological descriptions describe this process as the *stop* becoming affricated, synchronic phonetic evidence (illustrated below) suggests that it is the initial part of the close *vowel* or *glide* that becomes (devoiced and) fricated immediately after the stop release. Frication cannot be attributed merely to the narrower aperture of the close vowel or glide immediately after the stop release because (af)frication is not found in words such as *canyon*, *million*, *savior*, or *burial*, where the same glide or vowel apertures are involved but there are different preceding consonants. As there is no considerable pressure build up during the production of C1 in /nj, lj, vj, rj/ sequences, the air velocity is not high enough to cause

frication noise. Thus, it is the joint effect of a vowel or glide with a narrow aperture and the pressure build up for the stop which are responsible for the resulting frication.

Most likely, the same principles are responsible for the historical fricativization of high front vowels following sibilants in the transition from Middle Chinese to Modern Mandarin, for example, */si/ → [sʐ] ‘poetry’, */ʃi/ → [ʃʐ] ‘lion’ (Chen 1976). The high pressure build-up for the voiceless sibilant is released through the narrow constriction for the high vowel, generating frication.

The affrication of stops followed by high vowels and glides (i.e., segments impeding the free flow of air) is at the origin of a number of cross-linguistic sound patterns and present-day synchronic variation. Some of these are illustrated in (5) and (6) (see Hall and Hamann 2003 for a typological study of alveolar assibilation in 45 languages):

(5) Affrication of alveolars/dentals

Historical

a. English /tj dj/ > /tʃ dʒ/

/tʃ/ *actual, nature, mature, picture*

/dʒ/ or /dj/ *residual, soldier, remedial*

b. Latin /tj dj/ > Old Catalan /ts dʒ/ > Mod. Catalan /s (d)ʒ/

Latin *petia* > *peça* [‘pesə] ‘piece’

Latin *diurnum* > *jorn* [ʒorn] ~ [dʒorn] ‘day’

c. Japanese /t, d/ > [tɕ, dʒ]/__ /i/; /t, d/ > [ts, dʒ]/__ /u/ (= [ɮ])

/tii/ > [tɕii] *chii* ‘social status’

/tja/ > [tɕa] [tɕa] *cha* ‘tea’

/di/ > [dʒ] [dʒiɾemma] ‘dilemma’

/katu/ [katsɯ] ‘win’ (pres.)

d. Ikalanga (Mathangwane 1996, cited in Ohala 1997a), stop frication before the high front vowel /i/, and distinctive aspiration before high vowels /i, u/, but not before the next lower vowels /ɪ u/. The high and high-mid vowels have now merged.

Proto Bantu */tima/ > Ik. /ts^hima/ ‘well’

BUT PB*/tima/ > Ik. /tima/ ‘heart’

Proto Bantu */tudi/ > Ik. /t^hudzi/ ‘shoulder’

BUT PB */tundu/ > Ik. /tundu/ ‘basket’

Synchronic

e. English

[tʃ] or [tʃ̥] *Tuesday, tune, got you*

[tʃr] *train, truck;*

[tʃi:] *tea*

f. Brazilian Portuguese (Albano 1999)⁶

/di/ *Gandhi* [gãndʒi]

/ti/ *internet* [internetʃi]

g. Italian dialects (Tuttle 1997)

Standard Italian /tj/ *tieni* ‘hold’(imperative), Venetian [tʃeŋ]

Standard Italian *alti* (plural -i) ‘tall’ (pl.), Ticino [altʃ], [ɛltʃ]

(6) Velar softening

Historical

[k] / [g] + [i ɪ e y j] > [tʃ̥ ts̥ ʃ̥ s̥] / [dʒ̥ dz̥ ʒ̥ z̥]

a. English

O.E. *ciele, cele* [k] > *chill* [tʃ̥] (but ‘cold’ [k])

O.E. *cirice* [k] > *church* [tʃ̥] (cf Scots ‘kirk’ [k])

Gk. *Gymnasion* [g] > *gymnasium* [dʒ̥]

b. Italian

Lat. *cena(m)* [k] ‘dinner’ > *cena* [tʃe:na]

Lat. *regia(m)* [g] ‘palace’ > *regia* [reddʒa]

c. Tai (Li 1977: 221)

⁶ Palatalization and affrication of dental stops is a geographical dialect marker in Brazilian Portuguese. Dental stops are palatalized and affricated when followed by a high vowel [i], which may be lexical or inserted after a word-final dental stop in borrowed words (as BP only allows /s, r, n, l/ syllable-finally).

Lungchow	Po-ai
<i>kjau</i>	tʃau 'head', 'knot of hair on top of the head'
<i>kjaa</i>	tʃaa 'rice seedlings'
<i>kjooy</i>	tʃooy 'drum'

Synchronic

d. Paduan affrication (Krämer 2004)

	Standard Italian	Paduan
<i>ghiaccio</i>	[$\text{g}^{\text{h}}\text{attʃo}$]	[dʒaso] 'ice'
<i>chiama</i>	[$\text{k}^{\text{h}}\text{ama}$]	[tʃama] 'he/she calls'

The 'fronting' of /k g/+i j e/ sequences into alveolar or palatal affricates or fricatives has been the focus of much investigation (Grammont 1933, Bhat 1978, Guion 1998, to name a few). In contrast to articulatory accounts of the sound change, in terms of coarticulatory palatalization of the velar, Ohala (1989, 1992) provides an acoustic-auditory motivation for velar fronting. Chang, Plauché and Ohala (2001) present evidence that if the characteristic mid-frequency spectral peak of the burst in [ki] is degraded (and consequently perceptually missed), an alveolar sequence [ti] is reported by listeners, in line with confusion studies (e.g., Winitz, Scheib and Reeds 1972). Moreover, the [ki] tokens with the mid-frequency peak filtered out received better $\sqrt{\text{tʃ}}$ goodness scores than unfiltered tokens. Since variation of the acoustic cues of the stop burst influenced the direction of the consonant confusion –and paralleled the direction of the sound change– they argue that acoustic-auditory factors underlie 'velar fronting' (see also Ohala 1985, 1993). See Ohala (1983a, 1997c) and Plauché, Delogu & Ohala (1997) for asymmetries in the direction of confusion patterns and sound change, i.e., $\text{ki} > \text{ti}$ but not the reverse.

A corollary to the generalization that stops tend to engender frication on adjacent glides and high vowels is *D*.

D. Voiceless stops plus high vowels, glides and sonorants tend to be affricated more often than voiced stops.

Another generalization that can be accounted by aerodynamic factors is that frication especially emerges after voiceless stops, though it may also emerge after voiced stops (Ohala 1976), as illustrated in (5) and (6) above. The higher incidence of affrication in voiceless than

voiced stops has been noted cross-linguistically (Bhat 1978 for velar stops; Hall and Hamann 2003 for dental-alveolar stops) and is illustrated in the English example (5.a) above. While historically the /tj/ sequence in *nature*, *actual*, and *picture* has been lexicalised as $\widehat{tʃ}$, in comparable voiced sequences, e.g., *soldier*, *medial*, *individual*, the /dj/ > $\widehat{dʒ}$ / change has not lexicalised as often, and these words may be pronounced either [dj] or $\widehat{dʒ}$. Similarly, in German the sequence /tj/ developed into \widehat{ts} whereas /dj/ did not affricate (e.g., *nation* [na'tsjo:n] vs *indianisch* [m'dja:nɪʃ] 'indian'). The effect of the voicing of the stop on 'stop assibilation' in Latin and Romance languages has been observed by a number of investigators. For example, Pope (1952: 129, 131) notes that, in Latin, \widehat{tsj} is attested for /tj/ in the 4th century (e.g., *iustitia* [tsj] 'justice') but this process did not affect /dj/. Affrication of /dj/ is not reported till Late Latin when palatalization and affrication of both /tj/ and /dj/ are attested. Hall and Hamann (2003) on the basis of observed assibilation of /tj/ and /dj/ sequences in 45 languages posit that 'voiced stops cannot undergo assimilation unless voiceless ones do', such that there is an implication relationship /dj/ assibilation \supset [tj] assibilation.

The differential effect of voicing in the stop was addressed in section 2.2.1. To recall, the vibrating vocal folds for voiced stops constitute a relatively high resistance to air flowing from the lungs, and allow less air pressure to build up behind the stop constriction vis-à-vis voiceless stops, with an open glottis and a large and unimpeded flow. Consequently, there is a lower oral pressure at the release of a voiced stop, and thus less turbulence is generated (by equations 2 and 3). Empirical data corroborate that a higher amount of airflow and a longer duration of the release phase is found in /tjV/ than in /djV/ sequences (Hamann and Velkov 2005).

Since affricates frequently derive from stops with a long noisy release, a corollary to the diminished glottal flow and lesser affrication for voiced vis-à-vis stops, is the observation that voiced affricates are less frequent in languages of the world than voiceless affricates (ratio 1:3; Maddieson 1984: 38-39)

In sum, the generalization that 'stops engender friction on adjacent glides, high vowels and sonorants' is moderated by voicing effects, with 'voiceless stops tending to be affricated more often than voiced stops'.

II. Voicing impairs obstruency

For the same reasons that increased glottal flow during voiceless sounds favors supraglottal turbulence, the reduced glottal flow due to voicing, impairs the high intensity noise (and abrupt spectral discontinuities) characteristic of obstruency.

E. Voiced fricatives are hard to make; if voicing is strong, there is a tendency to de-fricate; if frication is achieved, there is a tendency to devoice.

Voiced fricatives are relatively difficult to produce due to the antagonistic aerodynamic requirements for frication at the supraglottal constriction (high oral pressure) and voicing (low oral pressure) (Note that this does not apply to the glottal fricative, [ɦ], for which both turbulence and voicing are generated at the vocal folds). Voiced fricatives require a pressure difference (ΔP) across the oral constriction sufficient to generate turbulence. This implies high oral pressure. That same high oral pressure, however, tends to impair the transglottal flow required for voicing. Thus, voiced fricatives involve very finely tuned aerodynamic conditions so that a pressure drop is maintained across both the glottal and the supraglottal constrictions⁷ (Ohala, 1983b; Solé 2002b).

During the production of voiced fricatives, if voicing is present, the reduced transglottal flow tends to impair strong frication (as intensity of turbulence is proportional to rate of flow), and if strong frication is achieved, the high oral pressure will tend to impair vocal fold vibration. Thus, voiced fricatives tend to devoice or to defricate, as evidenced synchronically and diachronically in the patterns below.

E.1 Voiced fricatives are relatively rare cross-linguistically.

The difficulty to produce simultaneous voicing and frication is reflected in segment inventories. Overall, voicing contrasts in fricatives are much rarer than in plosives, and they are found only in about a third of the world's languages as compared to 60 percent for plosive voicing contrasts (some languages, however, have voiced fricatives without corresponding voiceless fricatives emerging from weakened stops or fortition of initial approximants; Maddieson 2005). Furthermore, considering languages that utilize voicing with one of the

⁷ Solé (2002b) estimated the allowable range of aerodynamic variation for voiced fricatives from aerodynamic data. For one of her subjects, she estimated a subglottal pressure (P_s) of 7.6 cmH₂O during fricative production. Since transglottal flow to maintain voicing requires a pressure drop across the glottis ($P_s - P_o$) of at least 1-2 cmH₂O (and higher values to initiate voicing, 2-3 cmH₂O), that leaves a P_o of approximately 5.6 cmH₂O. Generation of turbulence for voiced fricatives ceases when the transoral pressure drops to about 3cmH₂O (Ohala, Solé and Ying, 1998; Catford, 1977:124; Stevens, 1998:480), which means that P_o may vary between a rather narrow range of 5.6-3 cmH₂O in order to sustain voicing and frication.

obstruent types but not the other, the probability of vocal fold vibration being absent on fricatives is double that found for stops (Ohala 1983b: 201).

E.2. Voiced fricatives tend to defricate

For the same magnitude of the oral constriction, voiced fricatives have a lower intensity of friction than voiceless fricatives, which makes them more likely to be perceptually heard as frictionless continuants (e.g., glides, rhotics, approximants) or missed altogether. The reason is diminished airflow through the glottis for voiced vis-à-vis voiceless fricatives due to vocal fold vibration (i.e., increased glottal resistance), and the need to keep oral pressure low for voicing. In addition, voiced fricatives are known to be shorter than voiceless fricatives, thus they allow less time for air to accumulate behind the constriction and create a high pressure build-up. These mechanisms -- lowered rate and duration of transglottal flow -- are responsible for a lower oral pressure, and a lower intensity of noise vis-à-vis voiceless fricatives (by equations (2) and (3)). As a phonological consequence voiced fricatives resemble more closely the so-called “frictionless continuants” such as [j w ɹ], and, indeed, diachronically this is often their ultimate fate, as illustrated in (7a, b). Approximants or frictionless continuants are the common phonetic manifestation of /v ð/ in Danish, e.g., [mað] *mad*, ‘food’ (cf. OE *mete*, Swedish and Norwegian *mat*, Icelandic *matur*). In Spanish and Catalan, the medial voiced stops /b d g/ are ‘spirantized’ to [β ð ɣ]⁸, see (7c), these latter sounds being more adequately described as approximants rather than fricatives (Martínez Celdrán 1991, 2004; Romero 1995). In Spanish these approximants may even disappear in some cases, for example, in past participles ending in –ado, e.g., *hablado* [a'βlao] ‘talked’ (cases of ‘hypercorrection’, i.e., insertion of non-etymological [ð] in similar sequences e.g., [βaka'laðo] for *bacalao* ‘cod’, are common).

(7) Defrication of voiced fricatives⁹

- a. Gliding and vocalization. In Middle English the voiced velar fricative allophone of /g/ (a voiced stop in OE) became /w/ or /u/ and the palatal allophone either became /j/ ~ /i/ or was lost (Mossé 1952).

⁸ Spirantization of voiced stops may be a maneuver to lower oral pressure in order to facilitate voicing (Ohala 1983b)

⁹ It has been reported that voiced stops may become voiced approximants without an intermediate fricative stage (see, e.g., Villafana Dalcher 2006 for Florentine Italian). In such cases defrication of voiced fricatives may not be necessarily at work.

[ɣ] > [w]	[i] > [j], [i]
OE <i>swelgan</i> > ME <i>swolwen</i> ‘swallow’	OE <i>genog</i> > ME <i>inough</i> ‘enough’
OE <i>boga</i> > ME <i>bow</i> ‘bow’	OE <i>mægden</i> > ME <i>maiden</i> ‘maid’
OE <i>sorg</i> > ME <i>sorow</i> ‘sorrow’	OE <i>sægde</i> > ME <i>said</i>

b. S-rhotacism (Solé 1992)

Latin *cerasea* > Catalan *cirera* ‘cherry’

Pre-lit. Catalan *Tolosanu* > Catalan *Tolrà, Toldrà* (placename)

English *was - were* (< O.E. *wesan*)

lost - forlorn (< O.E. *forleosan*)

Yurak *fire* ‘nest’ cf. Finnish *pesä*

Yurak *kuro-* ‘to cough’ cf. Lappish *gossâ-*

c. Spirantization

Spanish *sabe* /sabe/ [‘saβe] ‘(s)he knows’

cada /kada/ [‘kaðə] ‘each’

pega /pega/ [‘peɣa] ‘(s)he hits’

E.3 Voiced fricatives tend to be weakened or lost earlier than voiceless fricatives.

Interestingly, historical data indicate that voiced fricatives tend to be weakened or lost earlier than voiceless fricatives. This is illustrated in fricative weakening¹⁰ in Gallo-Romance. Preconsonantal /s/ was voiced before a voiced consonant, and [z] was weakened (into vowel, glide or tap) and lost as early as the 11th century, whereas voiceless [s] was pronounced well into the 13th century. Thus, for example, in Old French /s/ weakening and loss is found earlier in *blâmer* < *blasmer* < Lat. **blastemare* ‘blame’ and *mêler* < *mesler*, *medler* [ðl] < Lat. *misculare* ‘meddle’ than in *fête* < *feste* < Latin *festā* ‘holiday’ and *epuzer* < *espozer* < Latin **sponsare* (Pope 1952: 151, 449). Further, the different fate of etymological /s/ in the English words in (8) shows that [z] but not [s] had been lost at the

¹⁰ The term fricative ‘weakening’ is used here to indicate attenuation of the high frequency noise which characterizes fricatives, due to gestural reduction or aerodynamic factors. Fricative loss is considered the endpoint of the weakening continuum, i.e., extreme attenuation leading to the segment becoming inaudible. In perceptual terms gradient attenuation of the friction noise may result in identification of a discrete segment (e.g., a frictionless continuant, a vowel, a tap, an assimilated segment, or /h/) or in the perceptual loss of the segment (i.e., deletion).

time of the Norman Conquest, when the words were borrowed into English (Pope 1952: 151).

(8) [z] + voiced C	[s] + voiceless C
dine < O.Fr. disner	feast < O.Fr. feste
hideous < O.Fr. hisdos	espouse < O.Fr. spuse
male < O.Fr. masle	esquire < O. Fr. esquier

The aerodynamic and acoustic differences for voiced as opposed to voiceless obstruents have phonological significance in a number of patterns below (2.2.2.F; 2.2.3.I)

2.2.2 Changes in magnitude of the constriction.

Intensity of turbulence is dependent on the shape and area of the constriction through which the air has to pass. In this section we will review sound patterns involving the generation or impairment of turbulence due to variations in the area of constriction. Such variations in the cross-dimensional area of the constriction may result from coarticulation with adjacent sounds or position in the syllable.

F. Lingual fricatives tend to weaken when followed by consonants involving conflicting tongue configurations

Lingual fricatives exhibit highly constrained articulatory, aerodynamic and time requirements (Bladon and Nolan 1977). They require articulatory positioning to form a constriction within a certain critical range (approximately 0.1cm²; Stevens 1998: 47) and creating sufficient pressure difference across the oral constriction to generate frication. This requires sufficient rate of flow through the glottis and sufficient time to build up oral pressure behind the oral constriction. Precisely because they are highly constrained, lingual fricatives allow less articulatory and aerodynamic variation – in magnitude and time – than other segment types¹¹. Solé (2002a, b) has shown that apical trills have similar, if not more

¹¹ For example, because the constriction shape and area of lingual fricatives is critical, they tolerate less coarticulation with neighboring sounds (Recasens, Pallarès & Fontdevila, 1997), they are less overlapped (Byrd 1996), and they show lesser articulatory reduction in magnitude (Byrd and Tan, 1996) than other segment types (e.g., stops); and because temporal factors are also critical, fricatives are less susceptible to temporal reduction vis-à-vis other segments (Klatt, 1976; Byrd and Tan 1996).

constrained positional, shape, aerodynamic and elasticity requirements, and that they do not allow much articulatory variation if trilling is to be present.

FIG. 2

When lingual fricatives are followed by apical trills, involving antagonistic conflicting positional requirements of the tongue-tip/dorsum –raised and advanced tongue dorsum and a central groove for /s z ʒ ʒ/ vs. predorsum lowering and postdorsum retraction with a lax tongue-tip touching the alveolar ridge for the trill (see Figure 2)–, anticipatory tongue gestures for the trill may perturb the critical articulatory configuration (i.e., cross-sectional area of constriction) and/or temporal requirements for the generation of turbulence, and the fricative may be weakened or lost. Palatographic and aerodynamic evidence for fricative-trill sequences suggests that early onset of movements for the trill– within 30ms from the onset of lingual movements for the fricative – perturbs the articulatory trajectory and the critical constriction area for friction. In cases of lesser overlap, motor commands for the trill may arrive after the articulator attains the cross-sectional area for friction, but within the time needed to build up sufficient pressure difference to create audible friction (approximately 50ms from onset of oral pressure rise for voiced fricatives, with increased glottal resistance, and 30ms for voiceless fricatives. In such cases turbulent noise will not be generated (Solé 2002b).

S-weakening (into a tap or assimilated to the following sound) is also common before the conflicting lingual fricative /θ/ in Spanish (for example, /sθ/ > [rθ], [θ:] in *ascenso* ‘promotion’, *piscina* ‘swimming-pool’, Navarro Tomás 1980:111).

Examples of fricative weakening and loss in lingual fricative-trill sequences¹² are illustrated in (9).

(9) Examples of fricative weakening in lingual fricative-trill sequences

(a) Iberian Spanish

/sr/ *dos-reales* [ˌdoɾeˈales] ‘halfpenny’ (Navarro Tomás, 1980); *Osrám* [ˈoɾam]

/θr/ *voz ronca* [ˌboˈroŋka] ‘hoarse voice’, *Cruz Roja* [ˌkɾuˈroxa] ‘Red Cross’

¹² In reverse sequences, /rs/, where the trill is in coda rather than onset position, the trill is commonly detrilled and may assimilate to the fricative, e.g., Latin *bursa*, *morsicare* > Catalan *bossa*, *mossegar* ‘bag’, ‘to bite’ (Badia 1951: 202); /rs/ > [ʂ] (retroflex fricative) in Scots English (Bähr, 1974: 132ff) or Standard Swedish.

BUT:

/sl/, /sb/, /sk/ *desleal* [d̥ezle'al] 'disloyal', *esbozar* [ezβo'θar] 'to sketch', *asco* ['asko] 'disgust'

/θl/, /θm/, /θk/ *hazlo* ['aðlo] 'do it', *voz melodiosa* [boð melo'ðjosa] 'pleasant voice', *mezcla* ['meθkla] 'mixture'

(b) Catalan

/sr/ *les Rambles* [lə'rɑmbləs] (Recasens 1993)

/ʃr/ *mateix rotllo* [mə,t̪e(j)'rɔd̪lu] 'same story'

/ʒr/ *boig rematat* [bo(j)drəmə'tat̪] 'real crazy'

BUT:

/sl/, /sb/, /st/ *fes-li* ['fezli] 'do it' (for him), *les bledes* [ləz'βleðəs] 'the chard', *costellada* [kustə'ʎaðə] 'barbecue'

/ʃʎ/, /ʃp/ *mateix llit* [mə,t̪eʒ'ʎi t̪] 'same bed', *mateix poble* [mə,t̪eʃ'pɔplə] 'same town'

/ʒb/, /ʒp/ *boig valent* [boʒβə'len] 'brave crazy man', *boig per tu* [boʒpər'tu] 'mad about you'

(c) Portuguese

[ʒR] [ʒr] *dos reis* [du 'Rɛjʃ], [du 'rɛjʃ] 'of the kings', *Israel* [iRɐ'ɛʃ], [irɐ'ɛʃ]

Note that in the languages illustrated in (9) coda fricatives are voiced before a trill due to regressive voice assimilation, and thus they are more likely to be defricated (see E.2 above). Only the single trill realization is exemplified for the Spanish and Catalan data, but a long trill [r:] or a sequence [ɹr] ([ɹ] = fricative r) are also possible. The Portuguese data in (9c) illustrate that the uvular and apical variants of trills – the former involving the tongue dorsum – assimilate predorsal fricatives. In Italian, on the other hand, the fricative is preserved (e.g., /s # r/ *autobus rosso* [auɲobus^ə 'rɔsɔ] 'red bus', *Israele* [iz^dra'eɛ], [iz^dra'eɛ]) most probably due to the widespread insertion of epenthetic sounds at consonant release which allows the sequencing (*i.e.*, lack of overlap) of the gestures for the fricative and the trill. Whereas fricative to trill assimilation is probably an articulatorily gradient process (Solé 2002b), due to varying amounts of consonant overlap, the perceptual result is mostly categorical, that is, no frication is produced and the percept is commonly that of a trill or a long trill. Indeed, this process has led to reinterpretation in some placenames in Catalan, where the fricative has

disappeared in the lexical form, e.g., *Purroi* < etym. *Puigroig* /tʃ+r/ (Alcover and Moll, 1979), *Puigreig* [pu'retʃ] < [pudʒ'retʃ].

Additional evidence for the claim that the competing lingual requirements for trills impact on the generation of turbulence in preceding fricatives comes from (i) the more common and historically earlier weakening of lingual fricatives before trills (and also before laterals and nasals; see section 4) vis-à-vis other voiced consonants in Romance (Pope 1952:151footnote, 449; Rohlfs 1949; Torreblanca 1976; Recasens 2002: 352, 360), and (ii) electropalatographic and acoustic evidence that lingual fricatives lose their frication and are lost more commonly before trills (and also laterals and nasals) than before voiced stops and fricatives (e.g., in Majorcan Catalan, Recasens 2006).

G. Fricativization of syllable initial glides

Whereas the emergence of frication in glides in A.2 and C above was attributed to an increased rate of flow through the glottis when devoiced or when coarticulating with adjacent voiceless consonants, the frication of [j] word and syllable initially in dialects of Spanish, illustrated in (10a-c), is most likely due to a narrower oral constriction (principle 2). (Such narrowing of the constriction may give rise to an affricate [dʒ], see (10b, c)). There are two main reasons for this interpretation. One is that fortition (narrowing of a constriction) is common utterance and word-initially (Keating, Wright & Zhang 1999; Keating, Cho, Fougeron and Hsu 2003). Second, that fricativization of [j] also takes place in voiced contexts (intervocally in Argentinian Spanish, and after a voiced consonant in Iberian Spanish, as illustrated in (10b, c)) and, therefore, cannot be attributed to increased transglottal flow. It is the case, though, that these emergent fricatives may be devoiced in Argentinian Spanish, i.e., [ʃ]. Most likely, the devoiced variant results from increased narrowing of the constriction area and passive devoicing (see Ohala's 'aerodynamic voicing constraint' 1983b). Similar cases of fortition are found during the development of the Romance and Germanic languages.

The labial velar glide [w] is also fricativized or stopped word-initially in Spanish being pronounced [ɣw], [gw] and, less commonly, [βw] (see 10d).

(10) Frication of glides ([j] = voiced palatal fricative)

- a. *buey* [bweɣ] 'ox' vs *bueyes* ['bweɣes], ['bweɣes] 'oxen'

ley [lej] ‘law’ vs *leyes* ['lejes], ['lezes] ‘laws’

- b. Iberian Spanish (Navarro Tomás 1980:127ff.)

yo [jo], [jõ], [dʒo] ‘I’; *yeso* ['jeso], [ʎeso], ['dʒeso] ‘chalk’

cónyuge ['konjuxe], ['konʎuxe] ‘spouse’; *subyugar* [suβju'ɣar],

[subdʒu'ɣar] ‘subjugate’

- c. Argentinian Spanish (Colantoni 2006)

ayuda, [a'juða], [a'zuða], [a'dʒuða], [a'ʃuða] ‘help’

calle ['kaje], ['kaʒe], ['kadʒe], ['kafe] ‘street’

- d. Iberian Spanish (Navarro Tomás 1980:64)

huevo, ['weβo], [ʎweβo], [gweβo] ‘egg’

huelga, ['welya], [ʎwelya], [gwelya] ‘a strike’

2.2.3 Changes in oral pressure

Changes in oral pressure, such as those brought about by opening the velopharyngeal valve, will impact on the intensity of the resulting frication or stop burst. We review here a number of patterns illustrating that strong frication or a high intensity noise burst are difficult to achieve with concurrent or coarticulatory nasalization. Specifically, patterns showing that nasalization induces defricativization, and that buccal fricatives and stops do not tolerate nasalization. This leads to the generalisation that nasalization impairs obstruency.

III. Nasalization and obstruency do not mix.

Obstruents are characterized by high intensity frication and/or a noisy release burst. Since intense turbulence is pressure dependent (by equation (3b)), obstruents require a high build-up of air pressure behind the constriction in order to create audible turbulence when the pressure is released. If the obstruent constriction is downstream of the velopharyngeal port (i.e., ‘buccal’, labial to uvular), a tightly sealed velum is necessary to build up oral pressure. A lowered velum for nasality would vent the airflow through the nasal cavity, thus reducing or eliminating the required pressure difference across the oral constriction for intense turbulence. As a consequence, an open velopharyngeal port for nasality impairs high amplitude turbulence in buccal obstruents. In the case of glottal and pharyngeal fricatives and stops, for which the build up of pressure takes place further upstream than the velic

valve, a lowered velum would not affect the pressure build up, and thus they can be nasalized (Ohala 1975, Ohala and Ohala 1993). Nasalized glottal fricatives, /ɦ/, have been widely reported in languages (Ladefoged & Maddieson 1996: 131-134), and they occur phonetically in American English, e.g. *home* [hõõm]. Thus the requirement of a raised velum, and consequently the incompatibility of nasalization and obstruency, applies exclusively to obstruents articulated in front of the point of velic opening (buccal obstruents).

The antagonistic requirements of nasalization and obstruency, in addition to being predictable from aerodynamic and acoustic-auditory principles, have been demonstrated empirically. First, studies where oral pressure during the production of speech sounds was varied with a pseudo-velopharyngeal valve (a tube inserted at the side of the mouth via the buccal sulcus), simulating different degrees of nasalization (Ohala, Solé and Ying 1998), show that in producing a fricative there can be some opening of the velic valve, but the resistance at the velum has to be high relative to the resistance in the oral constriction so that the air will mostly escape through the aperture with lower resistance and create friction at the consonantal constriction. If resistance at the velopharyngeal port is lower than that at the oral constriction the air will escape through the nose (*i.e.*, the fricative will be nasalized), but supraglottal frication will be impaired. Ohala et al (1998) argue that velic openings which do not impair the build up of pressure for audible turbulence would be insufficient to create the percept of nasalization in the consonant or even adjacent vowels. Shosted (2006) obtained similar results with a mechanical model of the vocal tract with which he generated fricatives with different degrees of velopharyngeal opening.

Second, studies on velopharyngeal impairment (e.g., as presented in clinical cases of cleft palate) suggest that a velic opening of less than 10mm² during the production of oral stops exhibits normal aerodynamic values and can be tolerated without any perception of nasality, but velic openings of 10-20 mm² show diminished pressure and airflow values and the perception of inadequate nasal resonance (Warren, Dalston and Mayo 1993)¹³. Third, studies on coarticulatory nasalization in obstruents (e.g., Rothenberg 1968, Cohn 1990, Ohala and Ohala 1991; Basset, Amelot, Vaissière and Roubeau 2001) show that (mostly voiced) obstruents exhibit coarticulatory velic leakage preceding and following nasal vowels and nasal consonants, but that the velum may close before the release, allowing pressure to build up behind the constriction so as to produce frication or an oral burst, or that the velum may be slightly lowered throughout the obstruent, resulting in a relatively weak frication or burst.

¹³ This contrasts with the requirements for nasal consonants which require a velic opening greater than 20mm² and typically between 50 and 100mm².

The evidence presented so far suggests that obstruency and nasality do not mix. To the extent that an obstruent is a good obstruent perceptually (i.e., with intense frication or noisy release burst), it cannot be a good nasal (i.e., with perceptible nasal coupling); to the extent that it is perceptibly nasalized, it does not have the high amplitude noise cues for obstruency.

We review the phonological consequences of nasalization on fricatives and stops separately. We will address fricatives and nasalization first.

H. The rarity of nasal buccal fricatives.

Languages of the world have nasal stops, nasal taps, nasal approximants, nasal glides and nasal vowels but no nasal fricatives. Segments reported as nasalized fricatives are more adequately described as (i) frictionless continuants or approximants, due to the lack of high frequency aperiodic noise (e.g., in Umbundu, Schadeberg 1982; Coatzospan Mixtec, Gerfen 1996; and Waffa, Stringer and Hotz 1973) or (ii) as sequences of nasal and fricative segments, i.e., prenasalized fricatives (in Bantu languages, Kwa languages, and Igbo; Welmers 1973: 70-73) (Ohala 1983b, Ohala and Ohala 1993, Ohala et al. 1998). As shown above, if turbulence is created further upstream the point of velic opening – as in glottal and pharyngeal fricatives– velic lowering is of no consequence to the pressure build up, and the fricative can be simultaneously nasalized (see Ladefoged and Maddieson 1996:131-134 for examples).

I. Nasalization is associated with defricativization. The effect of voicing.

Nasalized fricatives, though rare, have been reported to occur in languages and it has been observed that they tend to be defricated if voiced –evidencing the difficulty to produce simultaneous frication and nasalization with reduced transglottal flow for voicing– and to lose their nasality if voiceless. For example, *voiced* nasalized fricatives are phonetically nasalized frictionless continuants (e.g., Waffa /β̃/ [β̃], Stringer and Hotz 1973; Umbundu /ṽ/ [ṽ], Schadeberg 1982), and voiced fricatives tend to lose their friction due to spreading nasalization and become nasalized approximants (e.g. [ṽ] ~ [ṽ ũ] in Guaraní, Gregores and Suarez 1967). In contrast, nasalized *voiceless* fricatives retain frication but do not differ much auditorily from non-nasalized fricatives, that is, the acoustic cues for nasalization are hardly detectable (Ohala 1975, Cohn 1993, Ladefoged and Maddieson 1996: 132).

The loss of friction in voiced but not voiceless nasalized fricatives follows from the aerodynamic factors reviewed. For the same degree of velopharyngeal opening, friction is more severely impaired in voiced than in voiceless fricatives. This is so because voiced nasalized fricatives have two additional mechanisms, other than nasal venting, impairing strong friction: (i) increased glottal resistance –which results in a lower oral pressure and inhibits the air vented through the nasal passage to be resupplied from the lungs (as it is the case for voiceless fricatives with an open glottis)– and (ii) the need to keep oral pressure low for voicing¹⁴. The differential effect of voicing in nasalized fricatives further illustrates the tendency for voicing to disfavor friction (section 2.2.1E). Other sound patterns illustrating the principle that nasalization induces defricativization are given in *J* and *K* below.

J. Failure of friction to emerge in a nasal context

Ohala (1983b:205-7) and Ohala and Ohala (1993: 228) provide the following examples of friction failing to emerge in a nasal context.

1. In the development from Middle Chinese to Mandarin, high vowels become fricatives when preceded by a sibilant fricative (e.g., */ʃi/ → [ʃ̺] ‘lion’), as stated in section 2.2.1C above. However, vowel assibilation fails to occur when the vowel is followed by a nasal consonant and is, consequently, nasalized. For example, */ʃiəm/ → [ʃən] ‘forest’ but not *[ʃ̺n] (Chen 1976).

2. In English, /h/ has the allophone [ç] before /j/, as in *huge* [çju:ɢ̺], as noted in section 2.2.1B above, but friction at the supraglottal constriction is not present if there is coarticulatory nasalization, e.g., *inhuman* [ĩn'h̃jũm̃ən], not *[ĩn'ç̃jũm̃ən].

3. In Yuchi, voiceless fricatives appear predictably between all vowels and following lingual stops, but fail to occur if the vowel is nasalized (Wagner 1934).

K. Fricatives are weakened or lost more often when followed by nasal than by non-nasal segments

¹⁴ Thus, Ohala, Solé and Ying (1998) report that when voiced and voiceless fricatives are vented with a pseudo-velopharyngeal valve –a tube inserted at the sides of the mouth via the buccal sulcus and the gap behind the molars– simulating different degrees of nasalization, when the valve has a similar impedance to that at the oral constriction (and as a result air is flowing out both through the nose and the mouth) voiced fricatives become frictionless continuants while voiceless fricatives retain their friction (though the intensity of friction is attenuated).

Coarticulation – the overlap of the articulatory configurations of contiguous segments – is well known and arises because it takes some minimum time to move articulators from one position to another. The antagonistic requirements of turbulence generation (a tightly closed velum to allow turbulent airflow in the oral tract) and nasal coupling (a lowered velum) in contiguous fricatives and nasals severely constrain the timing of velic movements if both segments are to be preserved. The relative phasing of velic and oral gestures in fricative + nasal (N or \tilde{V}) sequences have resulted in several sound changes, including (i) fricative weakening and loss, (ii) stop epenthesis, and (iii) vowel epenthesis (the latter two outcomes will be dealt with in section 4). Of interest here is fricative weakening or loss when followed by a nasal segment. Aerodynamic and acoustic data for fricative-nasal sequences shows that there might be anticipatory velopharyngeal opening for nasality during the acoustic duration of the fricative. Such nasal leakage diminishes the oral pressure build-up behind the fricative constriction, and attenuates the amplitude of frication, which may lead to fricative weakening or loss (Solé 2007a).

Although fricative weakening may also occur before non-nasals (e.g., Latin *misculare* ‘to mix’ > O.Fr. *mesler* > *mêler*; Germanic **bruzdon* ‘to embroider’ > Old Occitan *broidar* (where the ‘i’ is the result of the weakening process); Latin *festā* > French *fête* ‘holiday’) a number of scholars have noted that this process is favored by a following voiced consonant and, in particular, by a following [n], [m], [r] or [l] (Pope 1952: 151 footnote, 449, Rohlfs 1949, Torreblanca 1976, Recasens 2002: 352, 360). Whereas /s/ weakening before [r] may be attributed to antagonistic positional requirements of the tongue-tip and blade (see section 2.2.2.F), the weakening of fricatives before nasals (and laterals) may result from anticipatory velum (or tongue sides) lowering, thus affecting the aerodynamic requirements for the generation of turbulence.

Examples of fricative weakening due to coarticulatory nasalization resulting in vocalization or gliding (see 11a), rhotacism (exemplified in 11b), nasal assimilation (illustrated in 11c), and elision (see 11d) are found in historical sound change, morphophonological alternations and dialectal-stylistic variation.

(11) Examples of prenasal fricative weakening and loss

- a. [ɣn] > [jn], [wn] Latin *agnu* ‘lamb’, *ligna* ‘line’ > S. Italian dialects [‘ajənə], [‘lɛwna] (cited in Recasens 2002).
- b. [zn], [zm] > [rn], [rm] Latin **dis(ju)nare* ‘to eat breakfast’ > Old Occitan *dirnar/disnar* (cf. Cat. *dinar*) (Grandgent 1905: 53).

S. Spanish *mismo* ['mirmo] 'same' (Recasens 2002).

c. [zn] > [nŋ], [jn] *isn't* [inŋt], *ain't* [eɪnt]; *doesn't* [dʌnŋt]; *wasn't* [wɒnŋt]
(Gimson 1962).

BUT: *is there?* ['ɪzðə], *is she?* ['ɪʒi], *does he?* ['dʌzi]

d. [sm] [sn] > [m], [n] IE **gras-men* > Latin *grāmen* 'fodder', English *grama*,
gramineous;

BUT: IE **gras-ter* > Greek *gāster* 'stomach', English *gastric*, *epigastrium*
(Watkins 1985).

IE **dhus-no* > Welsh *dwn* 'dull, brown colour', OE *dun(n)* 'dark brown';

BUT: IE **dhus-ko* > Latin *fuscus*, OE *dox*, English *dusk* (Watkins, 1985).

L. Lower transitional frequency of fricatives followed by nasals

Fricatives combine less frequently with following nasals than with non-nasals. Solé (2007b) found a lower lexical frequency of word medial fricative + N sequences than of comparable fricative + C sequences in English, German and Dutch (the languages available in the CELEX database). Similarly, Rossato (2004) reports a bias against fricative + nasal sequences in a cross-linguistic count in 14 languages. Thus the transitional probabilities in the sequencing of sounds reflects the constraint against fricatives followed by nasal segments that endanger their high airflow requirements.

In sum, the data in *H* to *L* show that fricatives tend to lose their frication more often and earlier when they are nasalized, when they occur before a nasal vis-à-vis an oral sound, and that they combine less frequently with following nasals than with non-nasals, illustrating the generalisation that nasalization bleeds obstruency or, put another way, that fricatives do not tolerate nasalization.

M. In languages with nasal harmony, obstruents, including fricatives, block spreading nasalization

The incompatibility between obstruents and velic opening is evident in languages with nasal harmony. In such languages a nasal segment precipitates the spreading of nasalization to all following segments unless blocked by an oral obstruent. However, only buccal obstruents (labial to uvular), requiring a sealed velum, block spreading nasalization. The glottal obstruents [h ʔ], which do not require a raised velum since in their case frication is

